

POWER SUM SYMMETRIC FUNCTIONS

1. POWERSUM FAMILY

A powersum symmetric function with a single integer index, n , is the sum of all n^{th} powers of the variables $\{x_1, \dots, x_d\}$:

$$p_n = x_1^n + x_2^n + \dots + x_d^n$$

In terms of monomial symmetric functions,

$$p_n = m_{(n)}$$

The powersum symmetric function indexed by λ is the product of the corresponding single-indexed functions:

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots p_{\lambda_L} = p_1^{\rho_1} p_2^{\rho_2} \cdots p_n^{\rho_n} = p^\rho$$

in multi-index notation. Taken together $\{p_\lambda\}_\lambda$ form a basis for the symmetric functions under consideration.

The homogeneous and elementary symmetric functions have expansions in terms of the p_λ 's.

Definition Given $\rho = (1^{\rho_1} 2^{\rho_2} \cdots n^{\rho_n})$, define

$$z_\rho = 1^{\rho_1} 2^{\rho_2} \cdots n^{\rho_n} \rho_1! \rho_2! \cdots \rho_n!$$

with z_λ defined accordingly for $\rho = \rho(\lambda)$.

Proposition 1.1. *We have the expansions*

$$h_n = \sum_{\rho \vdash n} \frac{1}{z_\rho} p^\rho$$
$$e_n = \sum_{\rho \vdash n} \frac{(-1)^{n - \sum_j \rho_j}}{z_\rho} p^\rho$$

Example. With the partitions of 2, $\{[2], [11]\}$ and of 3, $\{[3], [21], [111]\}$, we have values of z_ρ for 2: (2,2) and for 3: (3,2,6) with

$$h_2 = p_2/2 + p_1^2/2$$

$$h_3 = p_3/3 + p_2p_1/2 + p_1^3/6$$

$$e_2 = -p_2/2 + p_1^2/2$$

$$e_3 = p_3/3 - p_2p_1/2 + p_1^3/6$$

etc.

1.1. Diagrams. The powersum symmetric functions correspond to a sum of single-rowed SSYT's with the same entry in each box. E.g., p_4 is the sum of SSYT's with shape



For $d = 3$, we have

$$p_4 =$$

$$\boxed{1\ 1\ 1\ 1} + \boxed{2\ 2\ 2\ 2} + \boxed{3\ 3\ 3\ 3}$$

the diagrams indicating the corresponding monomials. The number of terms is exactly d .